

# CadaFlex™

## SOFT CHEWS WITH MSM

CadaFlex™ is a comprehensive joint health supplement recommended to support joint function, flexibility, and healthy joint cartilage. It is formulated with our Proprietary Blend of Eggshell Membrane, Green-lipped Mussel Powder, Mushroom Powder, and Bacillus Coagulans, which contribute to digestive and immune health, promoting overall vitality.

### Available:

SP7702 For Small/Medium Dogs (84ct):

### Active Ingredients (per 4.8g soft chew):

Glucosamine HCl.....	600 mg
MSM.....	400 mg
Chondroitin Sulfate.....	250 mg
Turmeric.....	25 mg
Stratford Proprietary Blend (Egg shell Membrane, Green-lipped Mussel Powder, Maitake Mushroom Powder, Bacillus Coagulans).....	18 mg
Boswellia Serrata.....	10 mg
EPA (Eicosapentaenoic Acid) (fish Oil).....	.9 mg
DHA (Docosahexaenoic Acid) (fish oil).....	.6 mg

### Directions for use:

Weight	Initial Administration (4-6 Weeks)	Maintenance
Dogs under 10 lbs	1/2 soft chew per day	1/2 soft chew every other day
Dogs 10 to 29 lbs	1 soft chew per day	1/2 soft chew per day
Dogs 30 to 59 lbs	2 soft chews per day	1 soft chews per day

SP7700 For Large & Giant Dogs (84ct):

SP7701 For Large & Giant Dogs (150ct):

### Active Ingredients (per 7.5g soft chew):

Glucosamine HCl.....	900 mg
MSM.....	800 mg
Chondroitin Sulfate.....	350 mg
Turmeric.....	50 mg
Stratford Proprietary Blend (Egg shell Membrane, Green-lipped Mussel Powder, Maitake Mushroom Powder, Bacillus Coagulans).....	31 mg
Boswellia Serrata.....	.20 mg
EPA (Eicosapentaenoic Acid) (fish Oil).....	.12 mg
DHA (Docosahexaenoic Acid) (fish oil).....	.8 mg



### Directions for use:

Weight	Initial Administration (4-6 Weeks)	Maintenance
Dogs 60 lbs and over	2 soft chews per day	1 soft chew per day

**Inactive Ingredients:** Brewer's Yeast, Canola Oil, Glycerin, Hickory Smoke Flavor, Molasses, Pork Liver Powder, Potato Starch, Rice Bran, Sorbic Acid, Sunflower Lecithin, Tapioca Starch, and Vitamin E.

**Cautions:** If animal's condition worsens or does not improve, stop product administration and consult your veterinarian. Safe use in pregnant animals or animals intended for breeding has not been proven. Administer during or after the animal has eaten to reduce incidence of gastrointestinal upset. An examination from a veterinarian is recommended prior to using this product. May be irritant and not used in high doses with anticoagulant drugs or patients with stomach ulcers.

**Warning:** Not for human consumption. Keep out of the reach of children and animals. In case of accidental overdose, contact a health professional immediately.

